



PANDUAN PENGGUNA PADI

CADANGAN PENGGUNAAN



1 - 3 SEMBURAN SEMUSIM



KEKERAPAN: 30-35 hari lepas tanam, 50-55 hari lepas tanam, 70-75 hari lepas tanam

Tana-man	Kadar		Isipadu semburan L/ha	Kekerapan	Nota
	Per 10L	Per ha			
Padi	229 mL	8L	350 L	Setiap 2 minggu*	1 - 3 semburan semusim - Perkembangan tangkai - Sebelum berbunga - Selepas berbunga


*Bergantung kepada keadaan sawah

CARA PENGGUNAAN?



ARAHAN PENGGUNAAN

- Tambah air ke dalam tangki.
- Goncang bekas dengan kuat sebanyak 5 kali.
- Masukkan 1/3 botol ke dalam tangki bancuhan.
- Kacau larutan dan tuang ke dalam tangki sehingga habis.
- Jika ada mendapan putih di dasar bekas baja, masukkan 1/3 air ke dalam bekas dan kacau sehingga mendapan hilang, lalu campur ke dalam tangki bancuhan.
- Sedikit mendapan mungkin melekat pada bekas dan tidak memberi kesan kepada semburan.



USER'S GUIDE RICE

RECOMMENDATIONS FOR USE



1 - 3 TREATMENTS PER CROP CYCLE



FREQUENCY: 30-35 days after sowing, 50-55 days after sowing, 70-75 days after sowing

Crop	Rate		Spray volume L/ha	Frequency	Remarks
	Per 10L	Per ha			
Rice	229 mL	8L	350 L	Every 2 weeks*	1 - 3 applications/crop cycle - Panicle development - Early flowering - Late flowering

*Depending on the field conditions

HOW TO USE IT ?



INSTRUCTIONS FOR USE

- Add water to the tank.
- Shake vigorously by turning the can over 5 times.
- Empty 1/3 of the can into the tank.
- Repeat stirring and empty the rest into the tank.
- In case of white deposits at the bottom of the can after emptying, add 1/3 volume of water to the can, shake and stir into the solution.
- A small deposit can remain without impact on the use of the product.

